DECIGO: the Japanese Space Gravitational Wave Antenna

Masaki Ando[1,*], Seiji Kawamura[2], Takashi Nakamura[3], Naoki Seto[4], Kimio Tsubono[1], Kenji Numata[5], Ryuichi Takahashi[6], Mitsuru Musha[7], Ken-ichi Ueda[7], Ikkoh Funaki[8], Shigenori Moriwaki[9], Takeshi Takashima[8], Shin-ichiro Sakai[8], Takashi Sato[10], Nobuyuki Kanda[11], Shigeo Nagano[12], Mizuhiko Hosokawa[12], Takehiko Ishikawa[13], Shuichi Sato[2], Yoichi Aso[1], Mutsuko Y. Morimoto[2], Kazuhiro Agatsuma[14], Tomomi Akutsu[14], Tomotada Akutsu[5]6, Koh-suke Aoyanagi[15], Koji Arai[2], Yuta Arase[14], Akito Araya[16], Hideki Asada[17], Takeshi Chiba[18], Toshikazu Ebisuzaki[19], Motohiro Enoki[20], Yoshiharu Eriguchi[21], Feng-Lei Hong[30], Masa-Katsu Fujimoto[2], Mitsuhiro Fukushima[22], Toshifumi Futamase[23], Katsuhiko Ganzu[3], Tomohiro Harada[24], Tatsuaki Hashimoto[8], Kazuhiro Hayama[25], Wataru Hikida[26], Yoshiaki Himemoto[27], Hisashi Hirabayashi[8], Takashi Hiramatsu[27], Hideyuki Horisawa[28], Kiyotomo Ichiki[6], Takeshi Ikegami[30], Kaiki T. Inoue[31], Kunihito Ioka[3], Koji Ishidoshiro[1], Hiroyuki Ito[12], Yousuke Itoh[32], Shogo Kamagasako[14], Nobuki Kawashima[31], Fumiko Kawazoe[33], Hiroyuki Kirihara[14], Naoko Kishimoto[8], Kenta Kiuchi[15], Werner Klaus[12], Shiho Kobayashi[34], Kazunori Kohri[35], Hiroyuki Koizumi[36], Yasufumi Kojima[37], Keiko Kokeyama[33], Wataru Kokuyama[1], Kei Kotake[15], Yoshihide Kozai[38], Hideaki Kudoh[27], Hiroo Kunimori[12], Hitoshi Kuninaka[8], Kazuaki Kuroda[14], Kei-ichi Maeda[15], Hideo Matsuhara[13], Yasushi Mino[39], Jun-ichi Miura[7], Osamu Miyakawa[40], Shinji Miyoki[14], Tomoko Morioka[33], Toshiyuki Morisawa[26], Shinji Mukohyama[27], Isao Naito[41], Noriyasu Nakagawa[14], Kouji Nakamura[6], Hiroyuki Nakano[1], Kenichi Nakao[11], Shinichi Nakasuka[36], Yoshinori Nakayama[42], Erina Nishida[33], Kazutaka Nishiyama[8], Atsushi Nishizawa[43], Yoshito Niwa[43], Masatake Ohashi[14], Naoko Ohishi[44], Masashi Ohkawa[45], Akira Okutomi[14], Kouji Onozato[1], Kenichi Oohara[45], Norichika Sago[46], Motoyuki Saijo[47], Masaaki Sakagami[43], Shihori Sakata[33], Misao Sasaki[26], Masaru Shibata[21], Hisaaki Shinkai[48], Kentaro Somiya[49], Hajime Sotani[50], Naoshi Sugiyama[6], Hideyuki Tagoshi[46], Tadayuki Takahashi[8], Ryutaro Takahashi[2], Kakeru Takahashi[1], Hirotaka Takahashi[49], Takamori Akiteru[16], Tadashi Takano[8], Takahiro Tanaka[3], Keisuke Taniguchi[51], Atsushi Taruya[27], Hiroyuki Tashiro[3], Mitsuru Tokuda[11], Masao Tokunari[14], Morio Toyoshima[12], Shinji Tsujikawa[38], Yoshiki Tsunesada[52], Masayoshi Utashima[8], Hiroshi Yamakawa[53], Kazuhiro Yamamoto[14], Toshitaka Yamazaki[2], Jun'ichi Yokoyama[29], Chul-Moon Yoo[11], Shijun Yoshida[54], Taizoh Yoshino[55]

[1]*Department of Physics, The University of Tokyo, Bunkyo, Tokyo 113-0033, Japan,*
* *E-mail: ando@granite.phys.s.u-tokyo.ac.jp*
[2]*TAMA Project, National Astronomical Observatory of Japan, Mitaka, Tokyo 181-8588, Japan,*
[3]*Department of Physics, Kyoto University, Kyoto 606-8502, Japan,*
[4]*Department of Physics and Astronomy, University of California, Irvine, CA 92697-4575, U.S.A.,*
[5]*NASA Goddard Space Flight Center, Code 663, 8800 Greenbelt Rd., Greenbelt, MD20771, U.S.A.,*
[6]*Division of Theoretical Astronomy, National Astronomical Observatory of Japan, Mitaka, Tokyo 181-8588, Japan,*
[7]*Institute for Laser Science, The University of Electro-Communications, Chofu, Tokyo 182-8585, Japan,*
[8]*Institute of Space and Astronautical Science, Japan Aerospace Exploration Agency, Sagamihara, Kanagawa 229-8510, Japan,*
[9]*Department of Advanced Materials Science, The University of Tokyo, 5-1-5 Kashiwanoha, Kashiwa, Chiba 277-8561, Japan,*
[10]*Department of Electrical and Electronic Engineering, Faculty of Engineering, Niigata*

University, Niigata, Niigata 950-2181, Japan,
[11] *Department of Physics, Osaka City University, Osaka, Osaka 558-8585, Japan,*
[12] *National Institute of Information and Communications Technology (NICT), Koganei, Tokyo 184-8795, Japan,*
[13] *Institute of Space and Astronautical Science, Japan Aerospace Exploration Agency, Tsukuba, Ibaraki, 305-8505, Japan,*
[14] *Institute for Cosmic Ray Research, The University of Tokyo, Kashiwa, Chiba 277-8582, Japan,*
[15] *Department of Physics, Science and Engineering, Waseda University, Shinjuku, Tokyo, 169-8555, Japan,*
[16] *Earthquake Research Institute, The University of Tokyo, Bunkyo, Tokyo 113-0032, Japan,*
[17] *Department of Earth and Environmental Sciences, Hirosaki University, Hirosaki, Aomori 036-8560, Japan,*
[18] *Department of Physics, College of Humanities and Sciences, Nihon University, Setagaya, Tokyo 156-8550, Japan,*
[19] *RIKEN, 2-1 Hirosawa Wako 351-0198, Japan,*
[20] *Astronomical Data Center, National Astronomical Observatory of Japan 2-21-1, Osawa, Mitaka, Tokyo 181-8588, Japan,*
[21] *Department of Earth Science and Astronomy, The University of Tokyo, Komaba, Meguro, Tokyo 153-8902, Japan,*
[22] *Advanced Technology Center, National Astronomical Observatory of Japan, Mitaka, Tokyo 181-8588, Japan,*
[23] *Astronomical Institute, Tohoku University, Sendai 980-8578, Japan,*
[24] *Department of Physics, Rikkyo University, Toshima, Tokyo 171-8501, Japan,*
[25] *80 Fort Brown, Brownsville 78520, Texas, U.S.A.,*
[26] *Yukawa Institute for Theoretical Physics, Kyoto University, Kyoto, Kyoto 606-8502, Japan,*
[27] *Theoretical Astrophysics Group, Department of Physics, The University of Tokyo, Bunkyo-ku, 113-0033, Japan,*
[28] *Department of Aeronautics and Astronautics, School of Engineering, Tokai University,*
[29] *Research Center for the Early Universe, The University of Tokyo, 7-3-1, Hongo, Bunkyo-ku, Tokyo, 113-0033, Japan,*
[30] *National Institute of Advanced Industrial Science and Technology (AIST), Tsukuba, Ibaragi 305-8563, Japan,*
[31] *Kinki University School of Science and Engineering, Higashi-Osaka, Osaka 577-8502, Japan,*
[32] *Physics Department, University of Wisconsin - Milwaukee, P.O. Box 413, 2200 E. Kenwood Blvd., Milwaukee, WI 53201-0413, U.S.A.,*
[33] *Ochanomizu University Graduate School of Humanities and Sciences, Bunkyo, Tokyo, 112-8610 Japan,*
[34] *Astrophysics Research Institute, Liverpool John Moores University, Twelve Quays House, Egerton Wharf, Birkenhead L41 1LD, UK,*
[35] *Harvard-Smithsonian Center for Astrophysics, Cambridge, MA 02138, U.S.A.,*
[36] *Department of Aeronautics and Astronautics, University of Tokyo, Hongo 7-3-1, Bunkyo-ku, Tokyo 113-8656, Japan,*
[37] *Hiroshima University, Graduate School of Science, Higashi-hiroshima, Hiroshima 739-8526, Japan,*
[38] *Gunma Astronomical Observatory, Agatsuma-gun, Gunma 377-0702, Japan,*
[39] *Theor. Astrophysics, California Institute of Technology, Pasadena, CA 91125, U.S.A.,*
[40] *LIGO Laboratory, California Institute of Technology, M/C 18-34, Pasadena, CA 91125, U.S.A.,*
[41] *Numakage, Saitama-shi, Saitama 336-0027 Japan,*
[42] *Department of Aerospace Engineering, National Defense Academy, 1-10-20, Hashirimizu, Yokosuka 239-8686, Japan,*
[43] *Faculty of Intergrated Human Studies, Kyoto University, Kyoto 606-8501, Japan,*
[44] *MIRA Project, National Astronomical Observatory of Japan, Mitaka, Tokyo 181-8588, Japan,*
[45] *Department of Biocybernetics, Faculty of Engineering, Niigata University, Niigata, Niigata*

950-2181, Japan,
[46]*Department of Earth and Space Science, Osaka University, Toyonaka, Osaka 560-0043, Japan,*
[47]*Highfield, Southampton SO17 1BJ, United Kingdom,*
[48]*Department of Information Science, Osaka Institute of Technology, Kitayama 1-79-1, Hirakata, Osaka 573-0196, Japan,*
[49]*Max-Planck-Institut für Gravitationsphysik, Albert-Einstein-Institut, Am Mühlenberg 1, D-14476 Golm bei Potsdam, Germany,*
[50]*Department of Physics, Section Astrophysics, Astronomy and Mechanics, Aristotle University of Thessaloniki, Thessaloniki 54124, GREECE,*
[51]*Department of Physics, University of Illinois at Urbana-Champaign, 1110 West Green Street, Loomis Laboratory of Physics, Urbana, IL 61801, U.S.A.,*
[52]*Graduate School of Science and Engineering / Physics, Tokyo Institute of Technology, Ookayama, Meguro-ku, Tokyo, 152-8550, Japan,*
[53]*Research Institute for Sustainable Humanosphere, Kyoto University, Gokasho, Uji, Kyoto 611-0011, Japan,*
[54]*Pure and Applied Physics, Science and Engineering, Waseda University, Shinjuku, Tokyo 169-8555, Japan,*
[55]*Nakamura-minami, Nerima, Tokyo 176-0025, Japan,*
[56]*Department of Astronomy, The University of Tokyo, Hongo 7-3-1, Bunkyo-ku, Tokyo 113-0033, Japan*

DECIGO (DECI-hertz interferometer Gravitational wave Observatory) is the future Japanese space gravitational wave antenna with observation band around 0.1 Hz. It aims at detecting gravitational waves from various kinds of sources, with sufficient sensitivity to establish the gravitational wave astronomy. In the pre-conceptual design, DECIGO is formed by three drag-free spacecraft, 1000 km apart from one another. The relative displacements between proof masses housed in these spacecraft are measured by Fabry-Perot interferometers. We plan to launch DECIGO in 2024 after research and development phase, including two milestone missions (DECIGO pathfinder and Pre-DECIGO) for verification of required technologies.

Keywords: DECIGO, Gravitational waves, Astronomy, Space Mission

1. DECIGO

DECIGO (DECI-hertz interferometer Gravitational wave Observatory) is the future Japanese space gravitational wave (GW) antenna,[1] with observation frequency band of around 0.1 Hz (Fig. 1). This frequency band is the gap region between LISA (Laser Interferometer Space Antenna)[2] and terrestrial detectors such as Advanced LIGO[3] and LCGT (Large-scale Cryogenic Gravitational-wave Telescope).[4] In addition, this band opens the possibility to observe GWs from cosmological distance, because it is free from the confusion noises, irresolvable GW signals, from too many white dwarf binaries in our Galaxy.

Main targets of DECIGO are GWs from binary inspirals of compact binaries, and from the early universe. DECIGO will have sufficient sensitivity to observe GWs from distant (redshift of $z \sim 1$) neutron-star binaries which are a few months to 5 years before merger. By resolving GW signals emitted from many (about 3×10^5) binaries in this range, we will obtain information of mass distribution of neutron-stars, and thus, on the theory of the evolution of massive stars and on the equation of state of high-density matters. Moreover, observing distant binaries, which play



Fig. 1. Pre-conceptual design of DECIGO (left) and its design sensitivity (right).

as precise clocks, it will be possible to measure the acceleration of the expansion of the universe from their redshift change.[1] As for black-hole binaries, DECIGO will observe GWs from coalescences of intermediate-mass ($10^3 M_\odot$) black hole binaries, which could reveal the mechanism of the formation of super-massive black holes in the center of galaxies. The extremely good sensitivity of DECIGO would enable us to detect GWs from the very early universe, which could provide important information to understand the beginning of the universe.

2. Pre-conceptual design of DECIGO

In the pre-conceptual design, DECIGO is formed by three drag-free spacecraft, 1000 km apart from one another. Relative displacements of the proof masses (mirrors) inside the spacecraft are measured by Fabry-Perot interferometers (See Fig. 1). We adopted the Fabry-Perot configuration because it provides a better best sensitivity at 0.1 Hz band than an optical transponder configuration which is adopted by LISA. Although the Fabry-Perot configuration with shorter arm length has the larger acceleration noises by laser radiation-pressure noise and practical force fluctuations than transponder configuration with long arm length does, these noises would be still slightly lower than the confusion noise by Galactic binaries.

The distance between spacecraft (Fabry-Perot cavity arm length) was chosen to be 1000 km. This arm length was chosen so as to be short enough to avoid refraction losses of laser power, and to form Fabry-Perot cavities, and yet so as to be long enough to ensure the high sensitivity for GW signals. The mirrors forming the cavities, which works as proof masses in spacecraft, have a diameter of 1 m, with moderate reflectivity to realize the cavity finesse of 10. The mass of mirror (about 100 kg) was simply chosen to be the largest we could fabricate and handle. The laser source of DECIGO will have an effective power of 10 W with a wavelength of 532 nm. The orbit and constellation of DECIGO is to be determined, considering the gravity disturbances by the sun and planets, durability of the thruster fuels, solar power supply, and the required angle resolution for the GW source, and so on.

3. Milestone missions for DECIGO

Long and intensive development phase will be required in order to realize DECIGO. We plan to launch DECIGO in 2024 after design (a pre-conceptual design, a conceptual design, a preliminary design, and finally a final design) and prototype tests with the help of research and development with table top experiments. We also have two milestone missions, DECIGO pathfinder (DPF) and Pre-DECIGO, before the launch of DECIGO. DPF will be one small satellite consists of two proof mass mirrors, which form a short Fabry-Perot cavity. The cavity length is measured by a stabilized laser source, and the mirrors are kept in the satellite with a drag-free control. The target of DPF will the technical demonstrations: a drag free control, laser stabilization in space, precise measurement with Fabry-Perot cavity, and mirror clump system used at the launch of the satellite. In addition, since DPF will have a modest sensitivity for GW events, we expect some scientific results with continuous observation at the DECIGO frequency band. The objectives and a conceptual design of Pre-DECIGO will be determined during the research and development phase of DECIGO.

4. Conclusions

We have started a serious investigation to realize DECIGO by determining the pre-conceptual design. Although hard efforts will be required before its launch, DECIGO will provide fruitful scientific results by opening a new astronomy with gravitational waves.

References

1. N. Seto, S. Kawamura, and T. Nakamura, Phys. Rev. Lett, 87 (2001) 221103, S. Kawamura et al., Class. Quantum Grav. 23 (2006) S125.
2. LISA: System and Technology Study Report, ESA document ESA-SCI (2000).
3. "*LIGO II Conceptual Project Book*", LIGO M990288-A-M (1999).
4. K. Kuroda, et al., Class. Quantum Grav. 19 (2002) 1237.